

LEPTON CP ASYMMETRY IN B FACTORIES: ϵ_B AND NEW PHYSICS

Zhi-zhong Xing

Physics Department, Nagoya University, Chikusa-ku, Nagoya 464-01, Japan

In this talk, we first make some brief comments on the phase-convention dependence of the CP -violating parameter ϵ_B . A simple framework is then presented for analyzing new physics contributions to B^0 - $\bar{B}^0$ mixing and their effects on the lepton CP asymmetry, a promising signal to be further searched for in the first-round experiments of B factories.

1 Introduction

The study of various possible CP -violating phenomena in weak B -meson decays, starting from the end of 1970's, becomes more intensive today – on the eve of KEK and SLAC B factories, which will provide a unique opportunity to test the Kobayashi-Maskawa (KM) mechanism of CP violation in the standard model (SM) and to discover possible new physics (NP) beyond the SM.

Phenomenologically three different types of CP -violating signals are expected in neutral B -meson transitions: (a) CP violation from B^0 - $\bar{B}^0$ mixing; (b) CP violation from direct b -quark decay; and (c) CP violation from the interplay of decay and mixing. Type (a) can be detected most appropriately from the decay-rate asymmetry between two semileptonic channels $B^0 \rightarrow l^+ \nu_l X^-$ and $\bar{B}^0 \rightarrow l^- \bar{\nu}_l X^+$, the so-called “lepton CP asymmetry” (denoted as $\mathcal{A}_{SL}$ below). Type (b) may involve large uncertainties associated with the evaluation of hadronic matrix elements and (or) penguin contributions. Type (c) should play a crucial role in determining the KM CP -violating phase and (or) the NP phase(s), as it is independent of hadron or penguin pollution to a good degree of accuracy in some decay modes¹.

Experimentally, however, only the lepton CP asymmetry $\mathcal{A}_{SL}$ has been searched for to date. An upper bound on this asymmetry was first set by the CLEO time-integrated measurement: $|\mathcal{A}_{SL}| < 0.18$ or $\text{Re}\epsilon_B < 0.045$ at the 90% confidence level^{2,3}, where ϵ_B is a CP -violating parameter of B^0 - $\bar{B}^0$ mixing defined like ϵ_K in the neutral kaon system. The OPAL time-dependent measurement has recently given a more accurate result⁴: $\text{Re}\epsilon_B = 0.002 \pm 0.007 \pm 0.003$ in a special convention used for ϵ_B , which is equivalent to $\mathcal{A}_{SL} = 0.008 \pm 0.028 \pm 0.012$. Both experiments are consistent with the standard model prediction, i.e., $|\mathcal{A}_{SL}| \sim 10^{-3}$ or smaller. It should be noted, nevertheless, the existence of NP in B^0 - $\bar{B}^0$ mixing is possible to enhance $\mathcal{A}_{SL}$ up to the percent level⁵, observable in the first-round experiments of a B -meson factory with

about 10^8 events of $B^0\bar{B}^0$ mesons⁶.

The purposes of this talk are: (1) to comment on the phase convention dependence of the parameter ϵ_B ; and (2) to present a simple framework for analyzing NP effect on $\mathcal{A}_{\text{SL}}$. We expect that the further measurement of $\mathcal{A}_{\text{SL}}$ and its correlation with other CP asymmetries may serve as a sensitive probe for NP in B^0 - $\bar{B}^0$ mixing.

2 Comments on ϵ_B

In some literature, the CP -violating parameter ϵ_B is defined to relate the flavor eigenstates $|B^0\rangle$ and $|\bar{B}^0\rangle$ to the mass eigenstates $|B_1\rangle$ and $|B_2\rangle$:

$$|B_{1,2}\rangle = \frac{1}{\sqrt{2(1+|\epsilon_B|^2)}} [(1+\epsilon_B)|B^0\rangle \pm (1-\epsilon_B)|\bar{B}^0\rangle] . \quad (1)$$

Certainly ϵ_B has the phase freedom arising from the bound states $|B^0\rangle$ and $|\bar{B}^0\rangle$. The reason is simply that $|B^0\rangle$ and $|\bar{B}^0\rangle$ are defined by strong interactions only, leaving the relative phase between them undetermined. The dependence of ϵ_B on the bound-state phase convention can be removed, if one adopts the definition to link the CP eigenstates $|B_+\rangle$ (CP -even) and $|B_-\rangle$ (CP -odd) with the mass eigenstates $|B_1\rangle$ and $|B_2\rangle$:

$$|B_{1,2}\rangle = \frac{1}{\sqrt{1+|\varepsilon_B|^2}} (|B_\pm\rangle + \varepsilon_B|B_\mp\rangle) . \quad (2)$$

Note that ε_B and ϵ_B are not identical⁷, unless we fix the CP transformation between $|B^0\rangle$ and $|\bar{B}^0\rangle$ as $(CP)|B^0\rangle = \pm|\bar{B}^0\rangle$. Therefore it is better to adopt the phase-convention independent parameter ε_B , other than ϵ_B , in phenomenological applications⁸. In the analyses of CLEO and OPAL measurements^{2,4}, however, the phase-convention dependent parameter ϵ_B has been used.

The more important point is that both ε_B and ϵ_B depend on the phase convention of the KM matrix for quark flavor mixings, although ε_B itself is independent of the bound-state phases of $|B^0\rangle$ and $|\bar{B}^0\rangle$. The KM phase convention comes from the freedom in defining the quark field phases, thus different parametrizations of the KM matrix may result in different values for ε_B and ϵ_B . In this sense, even ε_B is *not* a physically well-defined parameter.

In the CLEO and OPAL analyses^{2,4}, $\text{Im}\epsilon_B = 0$ or $|\text{Im}\epsilon_B| \ll 1$ has been assumed in an undemonstrative way^{9,10}. That is why these analyses can yield an upper bound on $\text{Re}\epsilon_B$ solely from the measurement of the lepton CP asymmetry. To see this point more clearly, let us take the CLEO constraint on $\text{Re}\epsilon_B$ into account. The dilepton CP asymmetry measured on the $\Upsilon(4S)$

resonance reads

$$\mathcal{A}_{\text{SL}}(\epsilon_B) = \frac{4\text{Re}\epsilon_B(1 + |\epsilon_B|^2)}{(1 + |\epsilon_B|^2)^2 + 4(\text{Re}\epsilon_B)^2}, \quad (3)$$

and $\mathcal{A}_{\text{SL}}(\epsilon_B) = \mathcal{A}_{\text{SL}}(\epsilon_B)$ holds. Only in the assumption $|\text{Im}\epsilon_B| \ll 1$ (and $|\text{Re}\epsilon_B| \ll 1$), one can get $\mathcal{A}_{\text{SL}}(\epsilon_B) \approx 4\text{Re}\epsilon_B$. Then $|\mathcal{A}_{\text{SL}}(\epsilon_B)| < 0.18$ leads to $|\text{Re}\epsilon_B| < 0.045$. If one takes $|\text{Im}\epsilon_B| \sim 1$, however, $\mathcal{A}_{\text{SL}}(\epsilon_B) \approx 2\text{Re}\epsilon_B$ appears¹¹ and a different upper bound on $\text{Re}\epsilon_B$ must turn out from the same measurement of $\mathcal{A}_{\text{SL}}(\epsilon_B)$. Indeed the condition $\text{Im}\epsilon_B = 0$ corresponds to a special parametrization of the KM matrix which has $\text{Im}V_{td} = \text{Im}V_{tb} = 0$ (e.g., the one proposed recently by Fritzsch and the author¹²). In contrast, the condition $|\text{Im}\epsilon_B| \sim O(1)$ may be satisfied in the “standard” parametrization or the Wolfenstein form¹⁰.

3 NP effect on $\mathcal{A}_{\text{SL}}$

Now we present a simple framework to analyze NP contributions to $\mathcal{A}_{\text{SL}}$ and other CP asymmetries via B^0 - $\bar{B}^0$ mixing. In terms of the off-diagonal elements of the 2×2 B^0 - $\bar{B}^0$ mixing Hamiltonian $\mathbf{M} - i\mathbf{\Gamma}/2$, $\mathcal{A}_{\text{SL}}$ can be written as

$$\mathcal{A}_{\text{SL}} = \text{Im} \left(\frac{\Gamma_{12}}{M_{12}} \right). \quad (4)$$

In most extensions of the SM, NP can significantly contribute to M_{12} . However, NP is not expected to significantly affect the direct B -meson decays via the tree-level W -mediated channels. Thus $\Gamma_{12} = \Gamma_{12}^{\text{SM}}$ holds as a good approximation, where⁵

$$\Gamma_{12}^{\text{SM}} = - \frac{G_F^2 B_B f_B^2 M_B m_b^2}{8\pi} [(\xi_u^*)^2 T_u + (\xi_c^*)^2 T_c + (\xi_t^*)^2 T_t] \quad (5)$$

with KM factors $\xi_i \equiv V_{ib}^* V_{id}$ ($i = u, c, t$) for three quark families and QCD correction factors $T_u \sim -T_c \sim 0.1$ and $T_t \sim 1$. In the presence of NP, M_{12} can be written as

$$M_{12} = M_{12}^{\text{SM}} + M_{12}^{\text{NP}}. \quad (6)$$

Rescaling three complex quantities in Eq. (6) by $|M_{12}| = \Delta M/2$, where $\Delta M = (0.464 \pm 0.018) \text{ ps}^{-1}$ has been measured³, we obtain a parametrized triangle in the complex plane (see Fig. 1). The correlation between the SM and NP parameters reads

$$R_{\text{NP}} = -R_{\text{SM}} \cos 2(\theta - \phi_1) \pm \sqrt{1 - R_{\text{SM}}^2 \sin^2 2(\theta - \phi_1)}, \quad (7)$$




Figure 1: Triangular relation of M_{12}^{SM} , M_{12}^{NP} and M_{12} (rescaled by $\Delta M/2$).

where R_{SM} can be calculated in the SM box-diagram approximation⁵:

$$R_{\text{SM}} = \frac{G_{\text{F}}^2 B_B f_B^2 M_B m_t^2}{6\pi^2 \Delta M} \eta_B F\left(\frac{m_t^2}{m_W^2}\right) |\xi_t|^2 \quad (8)$$

with $\eta_B \approx F(m_t^2/m_W^2) \approx 0.55$. We see that there exist two solutions for R_{NP} .

The lepton CP asymmetry $\mathcal{A}_{\text{SL}}$ turns out to be⁵:

$$\begin{aligned} \frac{\mathcal{A}_{\text{SL}}}{C_m} = & R_{\text{SM}} R_{\text{NP}} \left[\text{Im} \left(\frac{\xi_u}{|\xi_t|} \right)^2 T_u + \text{Im} \left(\frac{\xi_c}{|\xi_t|} \right)^2 T_c + \text{Im} \left(\frac{\xi_t}{|\xi_t|} \right)^2 T_t \right] \cos(2\theta) \\ & + R_{\text{SM}}^2 \left[\text{Im} \left(\frac{\xi_u}{\xi_t} \right)^2 T_u + \text{Im} \left(\frac{\xi_c}{\xi_t} \right)^2 T_c \right] + \\ & R_{\text{SM}} R_{\text{NP}} \left[\text{Re} \left(\frac{\xi_u}{|\xi_t|} \right)^2 T_u + \text{Re} \left(\frac{\xi_c}{|\xi_t|} \right)^2 T_c + \text{Re} \left(\frac{\xi_t}{|\xi_t|} \right)^2 T_t \right] \sin(2\theta), \end{aligned} \quad (9)$$

where $C_m \approx 1.3 \times 10^{-2}$. Clearly the second term of $\mathcal{A}_{\text{SL}}$ comes purely from M_{12}^{SM} itself and its magnitude is expected to be of $O(10^{-3})$ due to the absence of the T_t contribution. The first and third terms of $\mathcal{A}_{\text{SL}}$ arise from the interference between M_{12}^{SM} and M_{12}^{NP} ; but they depend on nonvanishing $\text{Im}(M_{12}^{\text{SM}})$ and $\text{Im}(M_{12}^{\text{NP}})$, respectively. For appropriate values of θ and ϕ_1 , magnitudes of both the first and third terms of $\mathcal{A}_{\text{SL}}$ may be at the percent level! To obtain $|\mathcal{A}_{\text{SL}}| \sim O(10^{-2})$, however, there should not be large cancellation between two dominant terms in Eq. (9).

The CP asymmetry in $B_d \rightarrow J/\psi K_S$ can be calculated in the same framework. We obtain

$$\mathcal{A}_{\psi K} = R_{\text{SM}} \sin(2\phi_1) + R_{\text{NP}} \sin(2\theta). \quad (10)$$

As R_{NP} , R_{SM} and ϕ_1 , θ are dependent on one another through Eq. (7), $|\mathcal{A}_{\psi K}| \leq 1$ is always guaranteed within the allowed parameter space.

Two interesting cases, corresponding to $\text{Im}M_{12}^{\text{SM}} = 0$, $\text{Im}M_{12}^{\text{NP}} \neq 0$ and $\text{Im}M_{12}^{\text{NP}} = 0$, $\text{Im}M_{12}^{\text{SM}} \neq 0$, have been numerically illustrated by Sanda and the author⁵. It is found that $\mathcal{A}_{\text{SL}}$ does have good chances to reach the percent level, and the correlation between $\mathcal{A}_{\text{SL}}$ and $\mathcal{A}_{\psi K}$ (as well as other CP asymmetries) does reflect the information from NP.

4 Summary

We have commented on the dependence of ϵ_B or ε_B on the KM phase convention. We emphasize that neither of them can prove much advantage in describing data of CP violation from B^0 - $\bar{B}^0$ mixing. Also a simple framework has been presented for the analysis of possible NP contributions to B^0 - $\bar{B}^0$ mixing and of their effects on $\mathcal{A}_{\text{SL}}$ and other CP asymmetries. If we are lucky, we should be able to detect the lepton CP asymmetry at the percent level in the first-round experiments of B factories.

Acknowledgments: I would like to thank Y.Y. Keum for his warm hospitality and APCTP for its generous support, which made my participation in this wonderful workshop realizable. I am also grateful to A.I. Sanda for sharing physical ideas with me in a recent paper⁵, whose results were partially presented here. Finally useful discussions with S. Pakvasa, A.I. Sanda, S.Y. Tsai and D.D. Wu are acknowledged.

References

1. I.I. Bigi and A.I. Sanda, Nucl. Phys. B **193**, 85 (1981); **281**, 41 (1987).
2. CLEO Collaboration, J. Bartelt *et al.*, Phys. Rev. Lett. **71**, 1680 (1993).
3. Particle Data Group, R.M. Barnett *et al.*, Phys. Rev. D **54**, 1 (1996).
4. OPAL Collaboration, K. Ackerstaff *et al.*, CERN-PPE/97-036 (1997).
5. A.I. Sanda and Z.Z. Xing, Phys. Rev. D **56**, 6866 (1997).
6. H. Yamamoto, Phys. Lett. B **401**, 91 (1997); Y. Sakai, talk given at the B Physics Mini-workshop, September, Izu, Shizuoka, Japan (1997).
7. M.C. Bañuls and J. Bernabéu, CERN-TH/97-216 or hep-ph/9710348.
8. See, e.g., S.Y. Tsai, Mod. Phys. Lett. A **11**, 2941 (1996).
9. I would like to thank Y. Nir for pointing out this fact to me.
10. Z.Z. Xing, hep-ex/9709005.
11. See, e.g., T. Brown and S. Pakvasa, Phys. Rev. D **31**, 1661 (1985).
12. H. Fritzsch and Z.Z. Xing, Phys. Lett. B **413**, 396 (1997) (hep-ph/9707215); CERN-TH/97-201 or hep-ph/9708366 (Phys. Rev. D in press).